



## **AMERICAN ACADEMY OF PEDIATRICS**

### **Section on Pediatric Dentistry and Oral Health**

#### **POLICY STATEMENT**

Organizational Principles to Guide and Define the Child Health Care System and/or Improve the Health of All Children

### **Preventive Oral Health Intervention for Pediatricians**

**ABSTRACT.** This policy is a compilation of current concepts and scientific evidence required to understand and implement practice-based preventive oral health programs designed to improve oral health outcomes for all children and especially children at significant risk of dental decay. Additionally, it reviews cariology and caries risk assessment and defines, through available evidence, appropriate recommendations for preventive oral health intervention by primary care pediatric practitioners.

*Key words: pediatric oral health prevention, oral health intervention.*

## **PURPOSE/INTRODUCTION**

### **Review of Circumstances Leading to Development of This Policy**

Oral health is an integral part of the overall health of children.<sup>1</sup> Dental caries is a common and chronic disease process with significant consequences. As health care professionals responsible for the overall health of children, pediatricians frequently confront morbidity associated with dental caries. Because caries is a nonclassic infectious process (arising from shifts in subpopulation ratios of established normal flora), pediatricians have an opportunity to prevent, intervene, and, in collaboration with dental colleagues, manage this disease.

## **Preventive Oral Health Intervention for Pediatricians, *continued***

### **Justification of Policy**

The prevalence of dental caries for the youngest of children has not decreased over the past decade, despite improvements for older children.<sup>2</sup> Data from the Medical Expenditure Panel Survey revealed that 89% of infants and 1-year-olds had office-based physician visits annually, compared with only 1.5% who had dental visits. Consequently, visits to physicians outnumbered visits to dentists at 250 to 1 for this age group.<sup>3</sup> Because the youngest of the pediatric patient population visit the pediatrician more than the dentist, it is critical that pediatricians be knowledgeable about dental caries, prevention of the disease, and interventions available to the pediatrician and the family.

### **Rationale for Format**

This policy statement is an effort to assist the primary care pediatric practitioner in addressing issues of dental caries and general oral health. The statement begins by building a knowledge base regarding the caries process that can serve as a foundation for understanding prevention and intervention strategies. After explaining the science of cariology, assessment of caries risk is described to assist the pediatrician in deciding which preventive and interventional strategies need to be used. Specific prevention and intervention strategies are then described and explained.

Additionally, the concept and importance of the dental home as well as strategies to improve the connection of the medical and dental homes are presented. Lastly, recommendations are provided to assist the pediatrician with implementation of the provided information.

## **BACKGROUND CONCEPTS**

### **Cariology**

The most common oral disease encountered in children is dental caries. Dental caries is a

## **Preventive Oral Health Intervention for Pediatricians, *continued***

nonclassic infectious disease<sup>4</sup> that results from an interaction between oral flora and dietary carbohydrates on the tooth surface. To adhere to tooth structure, oral flora utilize dietary sugars to create a sticky biofilm that is referred to as dental plaque. Dietary sugar can change the biochemical and microbiologic composition of dental plaque. In the presence of a high-carbohydrate diet, cariogenic organisms constitute a greater portion of the total bacterial population.<sup>5,6</sup> Acids produced by bacterial fermentation of carbohydrates reduce the pH of dental plaque to the point at which demineralization of the enamel occurs. The initial carious lesion appears as an opaque white spot on the enamel, and progressive demineralization results in cavitations of the teeth. Dental caries is a process, and loss of tooth structure (a dental cavity) is an end stage in the process.<sup>7</sup>

Human dental flora, generally regarded as qualitatively stable once established and site specific to human dentition, is believed to consist of more than 1000 different organisms, of which only a limited number are associated with dental caries.<sup>8</sup> *Streptococcus mutans* is most strongly associated with dental caries and is considered to be an indicator organism of a subpopulation of cariogenic organisms. *Streptococcus mutans*, like its related cariogenic cohorts, has the ability to adhere to enamel and is uniquely equipped to produce significant amounts of acid (acidogenic) and endure within that acidic environment (aciduric).

Dental flora adheres to the teeth by creating a tenacious and highly complex biofilm referred to as dental plaque. Dental plaque is capable of concentrating dietary sugars; therefore, the chronic consumption of sugary foods and liquids will continually recharge the plaque matrix, resulting in copious supplies of sugars within the plaque matrix. *Streptococcus mutans* and other cariogenic flora will then ferment available sugars, resulting in high levels of lactic acid, a decreased local pH (approximately 5.0), and demineralization of dental enamel (at an

## **Preventive Oral Health Intervention for Pediatricians, *continued***

approximate pH of  $\leq 5.5$ ). Because *S mutans* and its aciduric cohorts continue to thrive at low pH, the resulting environment selects against nonaciduric flora, creating a shift in the subpopulation ratio of benign to aciduric flora. As this process continues over multiple generations, aciduric organisms incur an up-regulation of virulence genes that allow them to thrive at even lower pH (4.0). Diet-mediated shifts in subpopulation ratios of dental flora are instigated by significant sugar intake, (environmentally selecting for carious organisms). Therefore, significant sugar intake is a driving cause of the caries process.

### **Preventive Strategies**

An understanding of normal dental flora serves as a foundation for the development of preventive strategies, with 2 important considerations. First, dental flora exists in a symbiosis with the human species. Second, only a small number of the organisms within dental flora cause caries. Therefore, our objective is not to eliminate all dental flora but to suppress the cariogenic bacteria within the flora.

Preventive strategies can be differentiated into 2 distinct categories. Primary prevention involves optimization of maternal dental flora before and during colonization of the oral flora of the infant (during eruption of the primary dentition). This invaluable mode of prevention provides an opportunity for a reduction in the mother's constitutionally virulent, aciduric flora and down-regulation of virulence genes within the aciduric flora, decreasing the child's risk of dental decay, and is the basis for first dental visit recommendations at 1 year or earlier made by various medical and dental organizations. This mode of prevention and its adjuncts are reviewed in detail in the policy statement from the American Academy of Pediatrics, "Oral Health Risk Assessment Timing and Establishment of the Dental Home."<sup>9</sup>

Secondary prevention is the continual and ongoing management of subpopulation ratios

## **Preventive Oral Health Intervention for Pediatricians, *continued***

of benign and aciduric flora within dental plaque. This mode of prevention consists of managing the balance between causative factors and protective factors and is critical to preventing and reversing the caries process. Secondary preventive strategies are hierarchical and currently consist of dietary counseling, oral hygiene instruction, and judicious administration of fluoride modalities. Therefore, although all preventive modalities are important, modification of diet is most important, followed by oral hygiene compliance and then administration of fluorides.

By controlling risk factors before disease occurs, the probability of preventing disease, both in the immediate future and the long term, is improved. Preventive strategies for this complex, chronic disease require a comprehensive and multifocal approach that begins with caries risk assessment.

### **Caries Risk Assessment**

Caries risk assessment, based on developmental, biologic, behavioral, and environmental factors, evaluates the probability of enamel demineralization exceeding enamel remineralization over time. The goal of risk assessment is to anticipate and prevent caries initiation before the first sign of disease. During the period of 1999-2002, 41% of US children 2 to 11 years of age had caries in primary teeth.<sup>2</sup> An earlier study noted that 25% of children 5 to 17 years of age had 80% of the carious permanent teeth in the total population.<sup>10</sup> Assessing each child's risk of caries and tailoring preventive strategies to specific risk factors are necessary for improving oral health in a cost-effective manner.

Caries-risk assessment is very much a work in progress. In a systematic review of literature regarding risk factors in primary teeth of children 6 years and younger, a paucity of studies of optimal (ie, longitudinal) design was noted.<sup>11</sup> A study evaluating the reliability of multiple risk indicators determined there is no consistent combination of risk variables that

## **Preventive Oral Health Intervention for Pediatricians, *continued***

provide a good predictor of caries risk when applied to different populations across different age groups.<sup>12</sup> This study concluded that the best predictor of caries in primary teeth was previous caries experience, followed by parents' education and socioeconomic status.<sup>12</sup> Although previous caries experience cannot be used as a risk indicator for the predentate or very young child, white spot lesions, as precursors to cavities, can be considered analogous to previous caries experience when assessing the risk of a very young patient. An analysis of National Health and Nutrition Examination Survey (NHANES) III data revealed that children from households with low income levels are more likely to experience caries and have higher levels of untreated caries than their counterparts from higher-income households.<sup>13</sup> Collectively, children enrolled in Special Supplemental Nutrition Program for Women, Infants, and Children (WIC) programs, Head Start, or Medicaid are at higher risk than are children in the general population.

Caries risk factors unique to infants and young children include perinatal considerations, establishment of oral flora and host-defense systems, susceptibility of newly erupted teeth, dietary transitioning from breast and bottle feedings to cups and solid foods, and establishment of childhood food preferences. Although preterm birth per se is not a risk factor, a child with low birth weight may require a special diet or have developmental enamel defects or disabilities that increase caries risk. Early acquisition of *S mutans* is a major risk factor for early childhood caries and future caries experience.<sup>14</sup> A reduction of the salivary level of *S mutans* in highly infected mothers can inhibit or delay colonization of their infants.<sup>15</sup> Although evidence suggests that children are most likely to develop caries if *S mutans* is acquired at an early age, this may be compensated in part by other factors, such as good oral hygiene and a noncariogenic diet.<sup>11</sup> High-risk dietary practices appear to be established early, probably by 12 months of age, and are maintained throughout early childhood.<sup>16</sup> In addition to the amount of sugar consumed,

## **Preventive Oral Health Intervention for Pediatricians, *continued***

frequency of intake is important.<sup>17</sup> Sugar consumption likely is a more significant factor for those without regular exposure to fluorides.<sup>18</sup> Children experiencing caries as infants and toddlers have a much greater probability of subsequent caries in both the primary and permanent dentitions.<sup>19</sup>

Early risk assessment targets infants and young children who traditionally have yet to establish a dental home. Unrecognized disease and delayed care can result in exacerbated problems, leading to more extensive, costly, and time-consuming care.

Risk assessment strategies most applicable for screening purposes include those that are acceptable to patients, reliable, inexpensive, and performed easily and efficiently and require limited equipment/supplies. The American Academy of Pediatric Dentistry (AAPD) has developed a caries-risk assessment tool (CAT) for use by dentists and primary care practitioners familiar with the clinical presentation of caries and factors related to caries initiation and progression (see [http://www.aapd.org/media/Policies\\_Guidelines/P\\_CariesRiskAssess.pdf](http://www.aapd.org/media/Policies_Guidelines/P_CariesRiskAssess.pdf)).<sup>20</sup> Radiographic assessment and microbiological testing have been included in the CAT but are not required. Additionally, the American Academy of Pediatrics has created *Oral Health Risk Assessment Training for Pediatricians and Other Child Health Professionals*, which provides a concise overview of the elements of risk assessment and triage for infants and young children (see <http://www.aap.org/commpeds/doch/oralhealth/screening.cfm>).<sup>21</sup>

The chronic, complex nature of caries requires that risk be reassessed periodically to detect changes in the child's behavioral, environmental, and general health conditions. All available data must be analyzed to determine the patient's caries risk profile. Periodic reassessment allows the practitioner to individualize preventive programs and optimize the frequency of recall and dental radiographic examinations.

## **Preventive Oral Health Intervention for Pediatricians, *continued***

### **SPECIFIC PREVENTIVE STRATEGIES**

#### **Dietary Counseling**

Dietary counseling for optimal oral health in children should be an essential part of general health counseling. The recent policy statement from the American Academy of Pediatrics on prevention of pediatric overweight and obesity highlighted concerns about health problems in overweight children, including cardiovascular, endocrine, and mental health problems, and the importance of promoting healthy eating behaviors. Consumption of juice and sugar-sweetened beverages has been linked to childhood obesity and caries development.<sup>22-25</sup>

Sugars are a critical factor in caries development. Caries risk is greatest if sugars are consumed at high frequency and are in a form that remains in the mouth for longer periods.<sup>26</sup> Sucrose is the most cariogenic sugar, because it can form glucan, which enables bacterial adhesion to teeth and limits diffusion and buffering of acids. Although starch-rich foods pose a low caries risk, mixtures of finely ground, heat-treated starch and sucrose (eg, cereals, potato or corn chips) are also cariogenic.<sup>27</sup>

Human milk by itself does not promote tooth decay.<sup>28</sup> However, breastfed infants are at risk of caries when they receive sugary liquids or eat foods with sugars and fermentable carbohydrates.<sup>26</sup>

Parents and caregivers should be counseled on the importance of reducing exposure to sugars in foods and drinks. To decrease the risk of dental caries and to ensure the best possible health and developmental outcomes, it is recommended that parents do the following:

- Breastfeed infants during the first year of life and beyond as is mutually desired.<sup>29</sup>
- After nursing, remove the breast from a sleeping baby's mouth and cleanse the gums and teeth after feedings as well as before bedtime.



## **Preventive Oral Health Intervention for Pediatricians, *continued***

- Discourage a child's sleeping with a bottle. Any bottle taken to bed should contain only water.
- Limit sugary foods and drinks to mealtimes.
- Avoid carbonated beverages and juice drinks, (juice drinks contain high fructose corn syrup and <100% natural juice).
- Encourage children to drink only water and milk between meals.
- Encourage children to eat fruits.
- Limit the intake of 100% fruit juice to no more than 4 ounces per day.
- Foster eating patterns consistent with MyPyramid from the US Department of Agriculture.<sup>30</sup>

### **Optimal Use of Fluorides**

Fluoride, a naturally occurring element, has been instrumental in the widespread decrease in dental caries.<sup>31,32</sup> The mechanisms of fluoride are both topical and systemic, with evidence pointing to a greater topical effect.<sup>33</sup> Fluoride reduces enamel dissolution while it encourages remineralization.<sup>34</sup> Antimicrobial effects of fluorides at low pH are also significant.<sup>35</sup>

The delivery of fluoride includes community-based, professionally applied, and self-administered modalities. Water fluoridation is a community-based intervention that optimizes the level of fluoride in drinking water, resulting in pre and posteruptive protection of the teeth.<sup>36</sup> Water fluoridation is a cost-effective means of preventing dental caries, with the lifetime cost per person equaling less than the cost of 1 dental restoration.<sup>37,38</sup> In short, fluoridated water is the cheapest and most effective way to deliver anticaries benefits to communities.

Professionally applied topical fluorides (PATFs) have their greatest effect preventing caries and must be applied at regular intervals.<sup>39</sup> PATFs include gel, foam, in-office rinse, and

## **Preventive Oral Health Intervention for Pediatricians, *continued***

varnish. PATFs are safe and efficacious, with varnishes having the advantage of adherence to the tooth surface, decreasing likelihood of ingestion, and increasing time of contact between the fluoride and tooth surface.<sup>37,39</sup> In the primary dentition, varnish effectiveness (measured by percent of caries reduction) ranges from 30% to 63.2%,<sup>40,41</sup> and an analysis of the number of fluoride varnish applications received resulted in a dose-response effect that was enhanced when coupled with counseling.<sup>42</sup> Finally, self-administered fluorides, including dietary fluoride supplementation and fluoridated toothpaste, have proven effective, providing low but protracted elevation of fluoride concentrations.<sup>35,43</sup> Caries reduction associated with self-administered fluoride supplementation ranges from 32% to 72% in the primary dentition.<sup>40</sup> In children and adolescents, fluoride toothpastes, mouth rinses, and gels reduce dental caries to a similar extent.<sup>44</sup>

The decision to use fluoride therapies must balance the risk of caries against the risk of enamel fluorosis (hypomineralization of the developing enamel caused by excess fluoride ingestion). Patients determined to be at increased risk of dental caries are candidates for more aggressive fluoride therapy utilization. Caries susceptibility and sources of dietary fluoride (eg, water supplies, beverages, prepared food, toothpaste) should be considered before recommending fluoride therapies.<sup>45-48</sup> Enamel fluorosis develops before tooth maturation and emergence, typically in children younger than 8 years.<sup>49</sup> The risk of enamel fluorosis is an esthetic concern, with very mild or mild forms most commonly observed in the general population.<sup>2,50</sup>

### **ANTICIPATORY GUIDANCE**

Anticipatory guidance is the process of providing practical, developmentally appropriate information about children's health to prepare parents for significant physical, emotional, and

## **Preventive Oral Health Intervention for Pediatricians, *continued***

psychological milestones.<sup>51</sup> Anticipatory guidance during well-child visits is an effective tool to educate parents about maintaining children's health. Mirroring the pediatric model, the American Academy of Pediatric Dentistry advocates oral health anticipatory guidance.<sup>52-55</sup> Anticipatory guidance focused on oral health disease should be an integral part of preventive pediatrics. Information concerning the impact of diet on dental health and counseling in regards to oral hygiene, nonnutritive oral habits, and dental safety should be shared with parents. Therefore, in addition to dietary counseling and optimizing fluoride exposure, anticipatory guidance for oral health includes:

1. Infant oral hygiene instruction. Teeth should be brushed at least twice daily with caregiver supervision and assistance for children. For children with elevated dental caries risk, consider using a pea-sized or the same amount of toothpaste as the child's fifth digit fingernail. Flossing should begin as soon as adjacent teeth are in contact and for surfaces where 2 teeth touch and they can no longer be cleansed with a toothbrush.
2. Counseling regarding nonnutritive oral habits. Use of pacifiers in the first year of life may prevent sudden infant death syndrome.<sup>56</sup> Sucking habits (eg, pacifiers or digits) of sufficient frequency, duration, and intensity may be associated with dentoalveolar deformations. Some changes persist past the cessation of the habit. Professional evaluation is indicated for nonnutritive sucking habits that continue beyond 3 years of age.<sup>53</sup>
3. Age-appropriate information regarding dental injury prevention. Parents should cover sharp corners of household furnishings at the level of walking toddlers, ensure use of car safety seats, and be aware of electrical cord risk for mouth injury. Properly fitted mouth guards are indicated for youths involved in sporting activities with risk of orofacial

## **Preventive Oral Health Intervention for Pediatricians, *continued***

injury.

Anticipatory guidance is valuable, because it emphasizes prevention of dental problems rather than surgical or restorative care. Anticipatory guidance and well-child visits during the first 2 years of life decrease the number of hospitalizations among poor and near-poor children irrespective of race and health status.<sup>57</sup> Oral health anticipatory guidance can reduce dental expenditures.<sup>58</sup> In light of this evidence, oral health anticipatory guidance should be integrated as a part of comprehensive counseling during well-child visits.<sup>59</sup>

### **INTERPROFESSIONAL COLLABORATION AND ESTABLISHMENT OF A DENTAL HOME**

To be successful in preventing dental disease, interventions must begin within the first year of life. Pediatricians are well positioned to initiate preventive oral health care by providing early assessment of risk, anticipatory guidance, and timely referral to establish a dental home. The American Academy of Pediatric Dentistry, the American Dental Association, and the American Association of Public Health Dentistry recommend that infants be scheduled for an initial oral examination within 6 months of the eruption of the first primary tooth but by no later than 12 months of age.

The pediatric community promotes the concept of a medical home to improve families' care utilization, seeking appropriate and preventive services with optimal compliance to recommendations. The concept of the dental home is based on this model and is intended to improve access to oral care. A dental home is the ongoing relationship between the dentist and the patient, inclusive of all aspects of oral health care delivered in a comprehensive, continuously accessible, coordinated, and family-centered way.<sup>52,60,61</sup> A dental home should be able to provide the following:

## **Preventive Oral Health Intervention for Pediatricians, *continued***

1. An accurate risk assessment for oral diseases and conditions;
2. An individualized preventive dental health program based on risk assessment;
3. Anticipatory guidance about growth and development issues (eg, maxillofacial and dento-alveolar development);
4. A plan for emergency dental trauma management;
5. Information regarding care of teeth and oral soft tissues;
6. Nutrition and dietary counseling;
7. Comprehensive oral health care in accordance with accepted guidelines and periodicity schedules for pediatric oral health; and
8. Referrals to dental specialists, such as endodontists, oral surgeons, orthodontists, and periodontists when care cannot be provided directly within the dental home.

Lack of access to dental care can be a barrier to establishment of a dental home. Because of the specialized training and expertise, the dentist provides an ideal dental home; however, when a dentist is not available, the pediatric medical provider should fulfill the dictates of preventive oral health care until a dentist can be accessed and a dental home can be established. Therefore, primary care pediatric practitioners are an integral community component in the overall effort in addressing oral health issues (eg, access to care, preventive intervention). With the continuing challenges of access to dentistry coupled with preschool children making many more visits to medical offices than to dental offices, primary care practitioners with oral health training report that they have provided preventive oral health services for their pediatric patients.<sup>51,52</sup> North Carolina primary care practitioners were able to integrate preventive dental services into their practices, increasing preventive services for young Medicaid children whose access to dentists is restricted (eg, geographically or because of nonparticipation of dentists).<sup>62</sup>

## **Preventive Oral Health Intervention for Pediatricians, *continued***

Often, the first step in timely establishment of a dental home is a referral from the physician. Although a report from the US Preventive Services Task Force on physicians' roles in preventing dental caries in preschool children found referral by a primary care practitioner only partially effective in increasing dental visits,<sup>40</sup> another study<sup>63</sup> reported that dentists were more likely to see young children referred by primary care practitioners.

Primary care practitioners are able to identify children in need of a referral to a dentist.<sup>64</sup> After 2 hours of training in infant oral health, primary care pediatric practitioners accurately identified children with cavities with good specificity (92%-100%) and sensitivity (87%-99%).<sup>40,63</sup> These results suggest that dental screening can be incorporated into a busy pediatrics practice and that primary care pediatric practitioners can contribute significantly to the overall oral health of young children by encouraging parents to enroll their children in a dental home as early as possible.

In summary, the ideal setting for administration of oral health care is the dental home. When there is no access to a dentist, the pediatric medical provider should consider administering risk-based preventive oral health measures until a dental home can be made available. With preparation, primary care practitioners are routinely able to screen accurately and provide oral health anticipatory guidance for children. Furthermore, they are ideally positioned to refer children to a dental home in a timely manner. Establishing collaborative relationships between physicians and dentists at the community level is essential to increasing access to dental care for all children and improving their oral and overall health.

### **RECOMMENDATIONS FOR PRIMARY CARE PEDIATRIC PRACTITIONERS**

1. An oral health risk assessment should be administered periodically to all children.
2. Oral health risk assessment training should be recommended for medical practitioners

## **Preventive Oral Health Intervention for Pediatricians, *continued***

who are in training programs and those who currently administer care to children.

3. Dietary counseling for optimal oral health should be an intrinsic component of general health counseling.
4. Anticipatory guidance for oral health should be an integral part of comprehensive patient counseling.
5. Administration of all fluoride modalities should be based on an individual's caries risk. Patients who have a high risk of caries are candidates for consideration of more intensive fluoride exposure after dietary counseling and oral hygiene instruction as compared with patients with a lower risk of caries (see Fig 1 and 2).
6. Supervised use of fluoride toothpaste is recommended for all children with teeth.
7. The application of fluoride varnish by the medical practitioner is appropriate for patients with significant risk of dental caries who are unable to establish a dental home.
8. Every child should have a dental home established by 1 year of age.
9. Collaborative relationships with local dentists should be established to optimize the availability of the dental home.

## Preventive Oral Health Intervention for Pediatricians, *continued*

Figure 1




Figure 2

### High Caries Risk Protocol:

- Appoint Patient at 1 month intervals X 3.
- Review dietary intake of sugars sources, (juices, etc..) at each appointment.
- Assess Oral Hygiene at each appointment, (Plaque/Inflammation).
- Review fluoride exposure and apply Fluoride varnish at each appointment if risk factors persist.
- At the third 1 month visit, if all risk factors are well managed: Reappoint at 3 months, review diet, hygiene, fluoride exposure and apply fluoride varnish. If risk factors are not controlled: Continue with 1 month recalls until risk factors are managed.
- At 3 month recall interval, if all risk factors are well managed: Reappoint every 6 months, review diet, hygiene, fluoride exposure and apply Fluoride varnish.



## **Preventive Oral Health Intervention for Pediatricians, *continued***

### **CONCLUSION**

Oral health is an integral part of the overall health and well-being of children. A pediatrician familiar with the science of dental caries, capable of assessing caries risk, comfortable with applying various strategies of prevention and intervention, and connected to dental resources can contribute considerably to the health of his or her patients. This policy statement, in conjunction with the oral health recommendations of the American Academy of Pediatrics *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 3<sup>rd</sup> edition,<sup>65</sup> serves as a resource for pediatricians and other clinicians to be knowledgeable about addressing dental caries. With dental caries being such a common and consequential disease process in the pediatric population, it is essential that pediatricians include oral health in the daily practice of pediatrics.

**Preventive Oral Health Intervention for Pediatricians, *continued***

**SECTION ON PEDIATRIC DENTISTRY AND ORAL HEALTH EXECUTIVE COMMITTEE, 2006-2007**

Martha Ann Keels, DDS, PhD, Chairperson

\*Kevin J. Hale, DDS

Huw F. Thomas, DDS, MS, PhD

Martin J. Davis, DDS

Charles S. Czerepak, DMD, MS

Paul A. Weiss, DDS, Immediate Past Chairperson

**LIAISON**

Jack W. Morrow, DDS, MSD

American Academy of Pediatric Dentistry

**CONTRIBUTORS**

James Crall, DDS, ScD

David Krol, MD, MPH

Jessica Lee, DDS, MPH, PhD

Man Wai Ng, DDS, MPH

Rocio Quinonez, DDS

Jenny Stigers, DMD

**STAFF**

Aleksandra Stolic, MPH

*\*Lead author*

## Preventive Oral Health Intervention for Pediatricians, *continued*

### REFERENCES

1. National Institute of Dental and Craniofacial Research. *Oral Health in America: A Report of the Surgeon General*. Rockville, MD: US Department of Health and Human Services, National Institute of Dental and Craniofacial Research; 2000
2. Centers for Disease Control and Prevention. Surveillance for dental caries, dental sealants, tooth retention, endentulism, and enamel fluorosis—United States, 1988-1994 and 1999-2002. *MMWR Surveill Summ*. 2005;54(SS-3):1-43
3. Profile of Pediatric Visits (Tables 9-10), Based on 2000-2005 Medical Expenditure Panel Survey and 2000-2004 National Ambulatory Medical Care Survey. Elk Grove Village, IL: American Academy of Pediatrics. Updated June 2008. Available at: <http://practice.aap.org/public/ProfileOfPediatricVisits.pdf>. Accessed July 31, 2008
4. Fejerskov O. Changing paradigms in concepts on dental caries: consequences for oral health care. *Caries Res*. 2004;38:182-191
5. Berkowitz RJ. Etiology of nursing caries: a microbiologic perspective. *J Public Health Dent*. 1996;56:51-54
6. Bradshaw DJ, Marsh PD. Analysis of pH-driven disruption of oral microbial communities on vitro. *Caries Res*. 1998;32:456-462
7. Featherstone JDB. Prevention and reversal of dental caries: role of low-level fluoride. *Community Dent Oral Epidemiol*. 1999;27:31-40
8. Loesch WJ. Clinical and microbiological aspects of chemotherapeutic agents used according to the specific plaque hypothesis. *J Dent Res*. 1979;58:2404-2412
9. Hale KJ, American Academy of Pediatrics, Section on Pediatric Dentistry. Oral health risk assessment timing and establishment of the dental home. *Pediatrics*. 2003;111:1113-

## Preventive Oral Health Intervention for Pediatricians, *continued*

1116

10. Kaste LM, Selwitz RH, Oldakowski RJ, Brunelle JA, Winn DM, Brown LJ. Coronal caries in the primary and permanent dentition of children and adolescents 1-17 years of age: United States, 1988-1991. *J Dent Res.* 1996;75:631-641
11. Harris R, Nicoll AD, Adair PM, Pine CM. Risk factors for dental caries in young children: a systematic review of the literature. *Community Dent Health.* 2004;21(1 Suppl):71-85
12. Zero D, Fontana M, Lennon AM. Clinical applications and outcomes of using indicators of risk in caries management. *J Dent Educ.* 2001;65:1126-1132
13. Vargas CM, Crall JJ, Schneider DA. Sociodemographic distribution of pediatric dental caries: NHANES III, 1988-1994. *J Am Dent Assoc.* 1998;129:1229-1238
14. Berkowitz RJ. Mutans streptococci: acquisition and transmission. *Pediatr Dent.* 2006;28:106-109
15. Kohler B, Bratthall D, Krasse B. Preventive measures in mothers influence the acquisition of *Streptococcus mutans* in their infants. *Arch Oral Biol.* 1983;28:225-231
16. Douglass JM. Response to Tinanoff and Palmer: dietary determinants of dental caries and dietary recommendations for preschool children. *J Public Health Dent.* 2000;60:207-209
17. Gustaffson BE, Quensel CE, Lanke LS, et al. The Vipeholm dental caries study: the effect of different levels of carbohydrate intake on caries activity in 436 individuals observed for five years. *Acta Odontol Scand.* 1954;11:232-264
18. Burt BA, Pai S. Sugar consumption and caries risk: a systematic review. *J Dent Ed.* 2001;65:1017-1023
19. Li Y, Wang W. Predicting caries in permanent teeth from caries in primary teeth: an

## Preventive Oral Health Intervention for Pediatricians, *continued*

- eight-year cohort study. *J Dent Res.* 2002;81:561-566
20. American Academy of Pediatric Dentistry, Council on Clinical Affairs. Policy on use of a caries-risk assessment tool (CAT) for infants, children, and adolescents. *Pediatr Dent.* 2005-2006;27(7 Reference Manual):25-27. Available at:  
[http://www.aapd.org/media/Policies\\_Guidelines/P\\_CariesRiskAssess.pdf](http://www.aapd.org/media/Policies_Guidelines/P_CariesRiskAssess.pdf). Accessed October 17, 2007
  21. American Academy of Pediatrics. *Oral Health Risk Assessment Training for Pediatricians and Other Child Health Professionals.* Elk Grove Village, IL: American Academy of Pediatrics; 2005. Available at:  
<http://www.aap.org/commpeds/doch/oralhealth/screening.cfm>. Accessed July 31, 2008
  22. Krebs NF, Jacobson MS, American Academy of Pediatrics, Committee on Nutrition. Prevention of pediatric overweight and obesity. *Pediatrics.* 2003;112:424-430
  23. Ludwig DS, Peterson KE, Gormaker SL. Relation between consumption of sugar-sweetened drinks and childhood obesity: a prospective, observational analysis. *Lancet.* 2001;357:505-508
  24. American Academy of Pediatrics, Committee on Nutrition. Use and misuse of fruit juice in pediatrics. *Pediatrics.* 2001;107:1210-1213
  25. Dennison BA, Fockwell HL, Baker SL. Excess fruit juice consumption by preschool-aged children is associated with short stature and obesity. *Pediatrics.* 1997;99:15-22
  26. Tinanoff M, Palmer CA. Dietary determinants of dental caries and dietary recommendations for preschool children. *J Public Health Dent.* 2000;60:197-206
  27. Moynihan P. The British Nutrition Foundation Oral Health Task Force report—issues relevant to dental health professionals. *Br Dent J.* 2000;188:308-312

## Preventive Oral Health Intervention for Pediatricians, *continued*

28. Erikson PR, Mazhari E. Investigation of the role of human breast milk in caries development. *Pediatr Dent*. 1999;21:86-90
29. American Academy of Pediatrics, Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics*. 2005;115:496-506
30. US Department of Agriculture Web site. MyPyramid. Available at: <http://www.mypyramid.gov>. Accessed September 26, 2007
31. Centers for Disease Control and Prevention. Ten great public health achievements: United States, 1900-1999. *MMWR Morb Mortal Wkly Rep*. 1999;48:241-243
32. Petersen PE, Lennon MA. Effective use of fluorides for the prevention of dental caries in the 21<sup>st</sup> century: the WHO approach. *Community Dent Oral Epidemiol*. 2004;32:319-321
33. Hellwig E, Lennon AM. Systemic versus topical fluoride. *Caries Res*. 2004;38:258-262
34. ten Cate JM, Featherstone JD. Mechanistic aspects of the interactions between fluoride and dental enamel. *Crit Rev Oral Biol Med*. 1991;2:283-296
35. Lynch RJ, Navada R, Walia R. Low-levels of fluoride in plaque and saliva and their effects on demineralisation and remineralisation of enamel; role of fluoride toothpastes. *Int Dent J*. 2004;54(5 Suppl 1):304-309
36. Singh KA, Spencer AJ. Relative effects of pre- and post-eruption water fluoride on caries experience by surface type of permanent first molars. *Community Dent Oral Epidemiol*. 2004;32:435-436
37. Centers for Disease Control and Prevention. Recommendations for using fluoride to prevent and control dental caries in the United States. *MMWR Recomm Rep*. 2001;50(RR-14):1-42
38. Griffin SO, Jones K, Tomar SL. An economic evaluation of community water

## Preventive Oral Health Intervention for Pediatricians, *continued*

- fluoridation. *J Public Health Dent.* 2001;61:78-86
39. Hawkins R, Locker D, Nobel J, Kay EJ. Prevention. Part 7: professionally applied topical fluorides for caries prevention. *Br Dent J.* 2003;195:313-317
40. Bader JD, Rozier GR, Lohr KN, Frame PS. Physicians' roles in preventing dental caries in preschool children. A summary of the evidence for the U.S. Preventive Services Task Force. *Am J Prev Med.* 2004;26:315-325
41. Marinho VC, Higgins JP, Logan S, Sheiham A. Fluoride varnishes for preventing dental caries in children and adolescents. *Cochrane Database Syst Rev.* 2002;(3):CD002279
42. Weintraub JA, Ramos-Gomez F, Jue B, Shain S, Hoover CI, Featherstone JDB, Gansky SA. Fluoride varnish efficacy in preventing early childhood caries. *J Dent Res.* 2006;85(2):172-176
43. Marinho VC, Higgins JP, Logan S, Sheiham A. Topical fluoride (toothpastes, mouthrinses, gels or varnishes) for prevention dental caries in children and adolescents. *Cochrane Database Syst Rev.* 2003;(4):CD002782
44. Marinho VC, Higgins JP, Sheiham A, Logan S. One topical fluoride (toothpastes, mouthrinses, gels or varnishes) versus another for preventing dental caries in children and adolescents. *Cochrane Database Syst Rev.* 2004;(1):CD002780
45. Featherstone JD. The caries balance: the basis for caries management by risk assessment. *Oral Health Prev Dent.* 2004;2(Suppl 1):259-264
46. Jacobsen P, Young D. The use of topical fluoride to prevent or reverse dental caries. *Spec Care Dent.* 2003;23:177-179
47. Warren JJ, Levy SM. Systemic fluoride. Sources, amounts and effects of ingestion. *Dent Clin North Am.* 1999;43:695-711

## Preventive Oral Health Intervention for Pediatricians, *continued*

48. Levy SM, Kohout FJ, Kiritsy MC, Heillman JR, Wefel JS. Infant's fluoride ingestion from water, supplements and dentifrice. *JADA*. 1995;126:1625-1632
49. Pang DT, Vann WF Jr. The use of fluoride-containing toothpastes in young children: the scientific evidence for recommending a small quantity. *Pediatr Dent*. 1992;14:384-387
50. Riordan PJ. Perception of dental fluorosis. *J Dent Res*. 1993;72:1268-12674
51. Lewis CW, Grossman DC, Domoto PK, Deyo RA. The role of the pediatrician in the oral health of children: a national survey. *Pediatrics*. 2000;106(6):e84. Available at: <http://pediatrics.aappublications.org/cgi/content/full/106/6/e84>. Accessed September 26, 2007.
52. Nowak AJ, Casamassimo PS. Using anticipatory guidance to provide early dental intervention. *J Am Dent Assoc*. 1995;126:1156-1163
53. Nowak AJ, Warren JJ. Infant oral health and oral habits. *Pediatr Clin North Am*. 2000;47:1043-1066, vi
54. American Academy of Pediatric Dentistry. Guideline on periodicity of examination, preventive dental services, anticipatory guidance, and oral treatment for children. *Pediatr Dent*. 2005;27:84-86
55. American Academy of Pediatrics, Bright Futures Steering Committee. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. Hagen JF, Shaw JS, Duncan PM, eds. 3<sup>rd</sup> ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008
56. American Academy of Pediatrics, Task Force on Sudden Infant Death Syndrome. The changing concept of sudden infant death syndrome: diagnostic coding shifts, controversies regarding the sleeping environment, and new variables to consider in



## Preventive Oral Health Intervention for Pediatricians, *continued*

- reducing risk. *Pediatrics*. 2005;116:1245-1255
57. Hakim RB, Bye BV. Effectiveness of compliance with pediatric preventive care guidelines among Medicaid beneficiaries. *Pediatrics*. 2001;108:90-97
58. Savage MR, Lee JY, Kotch JB, Vann WF Jr. Early preventive dental visits: effects on subsequent utilization and costs. *Pediatrics*. 2004;114(4):e418-e423. Available at: <http://pediatrics.aappublications.org/cgi/content/full/114/4/e418>. Accessed September 26, 2007
59. Sanchez OM, Childers NK. Anticipatory guidance in infant oral health: rationale and recommendations. *Am Fam Physician*. 2000;61:115-120, 123-124
60. Poland C, Dean J. Pediatric oral health. In: Burns CE, Starr N, Brady MA, Dunn AM, eds. *A Handbook for Nurse Practitioners*. 2<sup>nd</sup> ed. Philadelphia, PA: WB Saunders Co; 2000
61. American Academy of Pediatric Dentistry, Council on Clinical Affairs. Policy on the dental home. *Pediatr Dent*. 2005;27:18-19
62. Rozier RG, Sutton BK, Bawden JW, Haupt K, Slade GD, King RS. Prevention of early childhood caries in North Carolina medical practices: implications for research and practice. *J Dent Ed*. 2003;67:876-885
63. Bentley EM, Holloway PJ. An evaluation of the role of health visitors in encouraging dental attendance of infants. *Community Dent Health*. 1993;10:243-249
64. Pierce KM, Rozier RG, Vann WF Jr. Accuracy of pediatric primary care providers' screening and referral for early childhood caries. *Pediatrics*. 2002;109(5):e82. Available at: <http://pediatrics.aappublications.org/cgi/content/full/109/5/e82>. Accessed September 26, 2007

**Preventive Oral Health Intervention for Pediatricians, *continued***

65. American Academy of Pediatrics, Bright Futures Steering Committee. Promoting oral health. In: Hagan JF, Shaw JS, Duncan PM, eds. *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*. 3<sup>rd</sup> ed. Elk Grove Village, IL: American Academy of Pediatrics; 2008:155-168

---

*All policy statements from the American Academy of Pediatrics automatically expire 5 years after publication unless reaffirmed, revised, or retired at or before that time.*